

**CE 5364 Groundwater Transport Phenomena  
Exercise Set 6**

**Exercises**

1. Improper waste disposal practices at an industrial site resulted in contamination of the soil on site by cadmium, a known carcinogen with a slope factor of  $6.10 \left(\frac{mg}{kgd}\right)^{-1}$ . We will consider the risk to off-site residents due to inhalation of airborne soil particles that include the cadmium. Based on monitoring data, the concentration of cadmium in the air off site is  $5.4 \times 10^{-4} \frac{mg}{m^3}$ .

Determine:

- (a) C<sub>Inh</sub> for residents that are children 1-6 years of age and adults.
- (b) The cancer risk due to these C<sub>Inh</sub> values for the children and adults.

Show all calculations and identify all parameter values used.

2. The same site also caused off-site lead concentrations that can cause non-cancer effects on the residents. The RfD for lead is  $6.90 \times 10^{-4} \left(\frac{mg}{kgd}\right)^{-1}$ . We will consider dermal exposures in this problem, with a lead concentration of  $260 \frac{mg}{kg}$  in the soil, and an absorption factor of 10 percent for the young children and 5 percent for adults.

Determine:

- (a) The NCDEX for residents that are children 1-6 years of age and adults.
- (b) The hazard quotients due to these NCDEX values for the children and adults.

Show all calculations and identify all parameter values used.

3. A contaminated groundwater that is a potential drinking water source has a manganese concentration of  $0.36 \frac{mg}{L}$ . The RfD for manganese is  $0.10 \frac{mg}{kg-d}$ . We will consider effects on children 6-12 (drinking 1 L/d) and adults (2 L/d).

Determine:

- (a) The NC<sub>Ing</sub> for children 6-12 and adults drinking this water.
- (b) The hazard quotients due to these NC<sub>Ing</sub> values for the children and adults.

Show all calculations and identify all parameter values used.

4. An animal exposure study was performed to determine an acceptable drinking water concentration for a chemical that causes liver disease in rats and is assumed to have a nonzero threshold. The following results were obtained.

**Control Group**

Comparison to historical records: no evidence of premature deaths Time of sacrifice: all surviving rats were sacrificed at 18 months Initial number: 100 Number of rats with liver disease: 3

**Test Group** Exposure conditions (lowest observed effect): 140 mg/L, 30 mL/d for a median of 12 months Time of sacrifice: all surviving rats were sacrificed at 18 months Comparison of weight and survival curves: no differences between test and control rats Median adult weight: 0.4 kg Initial number exposed: 100 Number of rats with liver disease: 12

Determine:

- (a) The LOAEL for the rats based on this study.
  - (b) The RfD for humans by adjusting for uncertainty. This result is subchronic animal data with no human exposure data available.
  - (c) Convert the RfD to an acceptable drinking water concentration.
5. Visit the EPA's IRIS system website (<http://www.epa.gov/iriswebp/iris/index.html>)

Determine:

- (a) Your favorite toxic or carcinogenic substance and print (or screen capture) the Quick View page for your choice.