

CE 5364 Groundwater Transport Phenomena
Exercise Set 2

Exercises

Most of these exercises will rely upon the materials in Asante-Duah, K. (1993) Hazardous Waste Risk Assessment. The necessary document excerpts are linked in the course notes.

1. Improper waste disposal practices at an industrial site resulted in contamination of the soil on site by cadmium, a known carcinogen with a slope factor of $6.10 \left(\frac{mg}{kgd}\right)^{-1}$. We will consider the risk to off-site residents due to inhalation of airborne soil particles that include the cadmium. Based on monitoring data, the concentration of cadmium in the air off site is $5.4 \times 10^{-4} \frac{mg}{m^3}$.

Determine:

- (a) C_{Inh} for residents that are children 1-6 years of age and adults.
- (b) The cancer risk due to these C_{Inh} values for the children and adults.

Show all calculations and identify all parameter values used.

2. The same site also caused off-site lead concentrations that can cause non-cancer effects on the residents. The RfD for lead is $6.90 \times 10^{-4} \left(\frac{mg}{kgd}\right)^{-1}$. We will consider dermal exposures in this problem, with a lead concentration of $260 \frac{mg}{kg}$ in the soil, and an absorption factor of 10 percent for the young children and 5 percent for adults.

Determine:

- (a) The NCDEX for residents that are children 1-6 years of age and adults.
- (b) The hazard quotients due to these NCDEX values for the children and adults.

Show all calculations and identify all parameter values used.

3. A contaminated groundwater that is a potential drinking water source has a manganese concentration of $0.36 \frac{mg}{L}$. The RfD for manganese is $0.10 \frac{mg}{kg \cdot d}$. We will consider effects on children 6-12 (drinking 1 L/d) and adults (2 L/d).

Determine:

- (a) The NC_{Ing} for children 6-12 and adults drinking this water.
- (b) The hazard quotients due to these NC_{Ing} values for the children and adults.

Show all calculations and identify all parameter values used.

4. An animal exposure study was performed to determine an acceptable drinking water concentration for a chemical that causes liver disease in rats and is assumed to have a nonzero threshold. The following results were obtained.

Control Group

Comparison to historical records: no evidence of premature deaths Time of sacrifice: all surviving rats were sacrificed at 18 months Initial number: 100 Number of rats with liver disease: 3

Test Group Exposure conditions (lowest observed effect): 140 mg/L, 30 mL/d for a median of 12 months Time of sacrifice: all surviving rats were sacrificed at 18 months Comparison of weight and survival curves: no differences between test and control rats Median adult weight: 0.4 kg Initial number exposed: 100 Number of rats with liver disease: 12

Determine:

- (a) The LOAEL for the rats based on this study.
 - (b) The RfD for humans by adjusting for uncertainty. This result is subchronic animal data with no human exposure data available.
 - (c) Convert the RfD to an acceptable drinking water concentration.
5. Visit the EPA's IRIS system website (<http://www.epa.gov/iriswebp/iris/index.html>)

Determine:

- (a) Your favorite toxic or carcinogenic substance and print (or screen capture) the Quick View page for your choice.